

High-precision astrometry on the VLT/FORS1 at time scales of few days [★]

P.F.Lazorenko¹, M.Mayor², M.Dominik³ **, F.Pepe², D.Segransan², and S.Udry²

¹ Main Astronomical Observatory, National Academy of Sciences of the Ukraine, Zabolotnogo 27, 03680 Kyiv-127, Ukraine

² Observatoire de Geneve, 51 Chemin des Maillettes, 1290 Sauverny, Switzerland

³ SUPA, University of St Andrews, School of Physics & Astronomy, North Haugh, St Andrews, KY16 9SS, United Kingdom

ABSTRACT

Context.

Aims. We investigate the accuracy of astrometric measurements with the VLT/FORS1 camera and consider potential applications.

Methods. The study is based on two-epoch (2000 and 2002/2003) frame series of observations of a selected Galactic Bulge sky region that were obtained with FORS1 during four consecutive nights each. Reductions were carried out with a novel technique that eliminates atmospheric image motion and does not require a distinction between targets and reference objects.

Results. The positional astrometric precision was found to be limited only by the accuracy of the determination of the star photocentre, which is typically 200–300 μ as per single measurement for bright unsaturated stars $B = 18 - 19$. Several statistical tests have shown that at time-scales of 1–4 nights the residual noise in measured positions is essentially a white noise with no systematic instrumental signature and no significant deviation from a Gaussian distribution. Some evidence of a good astrometric quality of the VLT for frames separated by two years has also been found.

Conclusions. Our data show that the VLT with FORS1/2 cameras can be effectively used for astrometric observations of planetary microlensing events and other applications where a high accuracy is required, that is expected to reach 30–40 μ as for a series of 50 frames (one hours with R_{special} filter).

Key words. astrometry – atmospheric effects – instrumentation: high angular resolution – planetary systems

1. Introduction

The provision of high-precision astrometry (of a few 100 μ as and below) is much desired, but so far it is not widely available. Over the recent years, photometric microlensing has proven its feasibility for not only detecting massive gas giant planets (Bond et al. 2004; Udalski et al. 2005), but even cool rocky/icy planets (Beaulieu et al. 2006). A fourth planet, of roughly Neptune-mass, was most recently claimed by Gould et al. (2006). Several further potential detections of planets by microlensing have however been missed (e.g. Jaroszyński & Paczyński 2002) since a proper characterization requires good data quality and a dense sampling. In fact, it is necessary to overcome ambiguities and degeneracies of binary lenses themselves (Dominik 1999), between mass ratio and source size (Gaudi 1997), and between binary-lens and binary-source systems (Gaudi 1998), as shown explicitly by Gaudi & Han (2004). The observation of the shift of the centroid of light composed of the lens star and the images of the source star with time, so-called astrometric microlensing (Høg et al. 1995; Hardy & Walker 1995; Miyamoto & Yoshii 1980; Dominik & Sahu 2000), provides a two-dimensional vector, whose absolute value is proportional to the angular Einstein radius (comprising information about the lens mass and relative lens-source parallax), in addition to the magnification provided by the photometric observations, where the latter only allows

to measure the angular Einstein radius if the finite size of the source star can be assessed in the light curve. While Safizadeh et al. (1999) first discussed the possibility of using astrometric microlensing for discovering and characterizing extra-solar planets, Han & Chunguk (2002) found that the planetary deviation to the astrometric signal expected from a Galactic bulge star lensed by a foreground star and a surrounding Jupiter-mass planet is about 10–200 μ as. Rather than requiring a stability of the detector on time-scales that correspond to the orbital period of planets as for the detecting of the astrometric shift seen in its host star due to motion around the common center-of-mass (e.g. Quirrenbach et al. 2004; Pravdo et al. 2005), the duration of the microlensing perturbations is just between a few hours and a few days.

Since ground-based monopupil telescopes were thought to be limited by atmospheric noise and therefore have rather poor precision (Lindgren 1980), only the use of high-precision optical interferometers such as SIM (Boden et al. 1998; Paczyński 1998), Palomar Testbed Interferometer (Lane & Mutterspach 2004), Keck (Colavita et al. 1998), and VLTI (Wittkowski et al. 2004) was considered so far for carrying out observations with such precision. A quick progress in adaptive optics, and its application to astrometry (Pravdo et al. 2005) leads to impressive results. However, the theoretical considerations by Lazorenko & Lazorenko (2004) and a first set of corresponding data (Lazorenko 2006, hereafter Paper I) demonstrated that astrometric measurements with very large ground-based telescopes are *not* atmospheric-limited. Instead, a high precision beyond this limit is achievable by elimination of atmospheric image motion at the data-reduction phase. For that purpose, a field of reference stars is configured into a virtual high-pass filter that

Send offprint requests to: P.F.Lazorenko

[★] Based on observations made with the European Southern Observatory telescopes obtained from the ESO/ST-ECF Science Archive Facility

** Royal Society University Research Fellow

absorbs most significant low-frequency modes of the image motion spectrum. For telescopes with a fully illuminated entrance pupil, the atmospheric error σ_{at} rapidly decreases with the increase of the telescope entrance pupil D . At $D \rightarrow \infty$, the asymptotic dependence is

$$\sigma_{\text{at}} = B_k R^{11/6} D^{-3/2} T^{-1/2} \quad (1)$$

where T is the integration time, R is the angular reference field radius, and B_k is the filter parameter. This relation involves an additional factor $\sqrt{R/D}$ as compared to that predicted by Lindegren (1980), so that large telescopes with $D = 8 - 10$ m and above are favourable. A specific amplitude apodization to the entrance pupil of future telescopes may further suppress the atmospheric error to $\sigma_{\text{at}} \sim R^{k/2} D^{-k/2+1/3} T^{-1/2}$ where k is the reduction parameter.

The astrometric performance of the VLT on very short time-scales could have been estimated from test reductions of a four-hour series of FORS2 frames obtained in the region of the Galactic bulge (Paper I). A positional accuracy of $300 \mu\text{s}$ per single $R=16$ mag star measurement at 17 sec exposure was shown to be limited only by the number of photons received from the observed star.

The current study extends the certification of high-accuracy astrometric performance with the VLT to time scales of 1–4 days, which is sufficient for monitoring astrometric planetary deviations in microlensing events. In Sect. 2 of this paper, we lay out the observational basis of this investigation, describe the image processing, and show how the stellar centroids are computed. Particular problems in astrometric reductions that are based on background reference stars with unstable positions, and a new reduction method are discussed in the subsequent Sect. 3. Reduction of the observational data with this method is described in Sect. 4. Sect. 5 focusses on the noise and correlations in the measurements that, if exist, can be related to the different parts of the telescope (optics, mechanics, electronics), to the atmosphere, etc., preventing statistical improvement of the accuracy by means of simple accumulation of the number of frames.

2. Observations and computation of centroids

For our study, we were able to use VLT frames that are available in the ESO Archive: those obtained with the FORS1 camera of UT1 in the sky region near the neutron star RX J0720.4-3125 under the ESO program 66.D-0286 (Motch et al. 2003) were found to be best-suited for our purposes. Images were acquired during 4 nights from 20 to 23 December 2000 (40 frames) and 4 nights from 29 December 2002 to 02 January 2003 (25 frames). During each 5 to 12 frames were obtained in B filter with ~ 10 mn exposure time and FWHM varying from 0.45 to 0.85'', where the high-resolution mode with 0.10''/px scale and about $3.3 \times 3.3'$ field-of-view was used.

These data enable us to carry out a detailed and comprehensive study of the astrometric precision that is achievable with the VLT/FORS1 at short time scales. Moreover, our analysis serves as a test reduction for future observations. Unfortunately, between some frames, the images turned out to be displaced by more than 200 px. For that reason, peripheral stars frequently needed to be omitted for much jittered frames. Moreover, for astrometric purposes, the choice of the B filter is not optimal, since it results in larger residual atmospheric chromatism as compared to R_{special} . However, it were just these unfortunate circumstances that forced us to look for a reduction that significantly improves the quality.

Observations were obtained with the Longitudinal Atmospheric Dispersion Corrector (LADC), which is a two-prism optical device (Avila et al. 1997) that allows to correct for atmospheric differential chromatic refraction (DCR). DCR is caused by the atmosphere, which acts like a prism with a certain dispersion and thereby smears and displaces stellar images (e.g. Monet et al. 1992). The size of this effect depends on the zenith distance and the colour of the observed star and can be modelled as shown in Eq. (12).

Raw images were debiased and flat-fielded using master calibration frames taken from the ESO Archives. Positions x, y of stellar centroids were computed using the profile fitting technique that had already been applied earlier for the high-precision astrometric reduction of FORS2 images (Paper I). Stellar profiles in 10×10 px windows were fit by a sum of three modified elliptic Gaussians involving 12 free model parameters altogether. The dominant Gaussian centered at x, y with extent parameters σ_G^x, σ_G^y along corresponding coordinate axis and further specified by its orientation angle, was considered to contain a flux I mounting to about 2/3 of the total light received from the star. Other auxiliary Gaussians allowed profile fitting to the photon noise limit at which residuals of pixel counts from the model are characterized by $\chi^2 \approx 1.0/\text{px}$ (except for central parts of images of bright stars). Objects with $\chi^2 \geq 3.0/\text{px}$ deviations were rejected and only unsaturated images with $B > 18$ mag were processed.

Due to the complex shape of the PSF, the precision ε of the image centroiding was obtained by numerical simulation, for which a set of randomly generated star images with profiles comparable to the observed ones was created and where Poisson noise was added to pixel counts. This simulation shows that ε can be well-approximated by

$$\varepsilon = \frac{\theta}{2.46 \sqrt{I}} (1 + \alpha_0 I^{-0.7}), \quad (2)$$

where θ denotes the seeing (which turned out to be $\theta = 3.10 \sigma_G$ in our three-component model), I refers to the electron count, and α_0 is a characteristic constant that depends on the background noise. For the model background fluxes b_0 of 810 or 540 e^-/px (average values at 2000 and 2002-2003 epochs), α_0 was found to be 1150 or 887, respectively. Eq. (2) becomes an exact relation only for a fixed average seeing $\theta_0 = 0.608''$ at which $\sigma_G = 1.96$ px, but it provides a reliable approximation for variable FWHM of that order.

3. The reduction process

3.1. Filtration of atmospheric image motion

Let us shortly recall the previously discussed process of attenuation of the image motion spectrum (Lazorenko & Lazorenko 2004), which is based on the virtual symmetrization of the reference field. Consider some region of the sky with N stars being imaged in $m = 0, 1 \dots M$ consecutive frames, where x_{lm}, y_{lm} are the centroids of $l = 1, 2 \dots N_m$ stars measured in frame m . For various reasons (jittered images, cosmics, outliered data, etc.) a significant fraction of centroids cannot be obtained from every frame, therefore $N_m \leq N$. Here and below, indices i, j, l refer to the target, reference, and star of any type, respectively, and $m = 0$ is assigned to the reference frame. Upper indexes x, y , if used, refer to the coordinate axes. A list of variables most frequently used in the Paper with a short description is given in Table 1.

The differential position X_{im} of the i -th target star in the m -th frame is obtained from CCD coordinate differences $x_{im} - x_{jm}$ of

Table 1. A list of frequently used variables

quantity	definition	first use, equation
x, y	measured star positions	(3)
X, Y	positions x, y filtered of image motion and given in the system of local reference stars with "fixed" positions	(3)
λ^x, λ^y	same as X, Y but given in a single all-frame system with "floating" reference stars	(9)
ξ^x, ξ^y	displacement (average for a given frame series) of the star from its position in reference frame, for astrometric reasons (proper motion etc.)	(14)
$\hat{\xi}^x, \hat{\xi}^y$	same as ξ^x, ξ^y but given with respect to the mathematical expectation of the star position in reference frame	(10)
ρ	chromatic coefficient; atmospheric displacement of the star image is proportional to ρ	(12)
d	chromatic coefficient; describes compensating action of the LADC	(12)
h	differential chromatic displacement (DCR effect) of the star image	(12)
$\mathcal{M}$	atmospheric image motion effect in x, y	(10)
$\Delta\mathcal{M}$	residual atmospheric image motion in X, Y	(14)
e	error of photocenter measurement	(10)
ε	variance of e	(1,25)
u	residuals of conditional equations (23)	(26)
$\mathcal{N}$	noise in observations that enters Eqs.(23)	(17)
σ^2	observed variance of $\mathcal{N}$ found from residuals u	(26)

the target and field star centroids. More specifically, it is given as the weighted average

$$X_{im} = \sum_{j=1}^N a_{ij}(x_{im} - x_{jm}) = x_{im} - \sum_{j=1}^N a_{ij}x_{jm}, \quad m = 1, 2 \dots M \quad (3)$$

of these differences, where the weights a_{ij} satisfy the normalization condition

$$\sum_{j=1}^N a_{ij} = 1 \quad (4)$$

and are chosen, so that the effect of image motion effect in X_{im} is minimized. The prime indicates that the summation is carried out over the subset Ω_{im} of stars used as reference for i -th star only, where the index $j = i$ is omitted. The corresponding positional y -coordinate Y_{im} is obtained in analogy to Eq. (3).

Atmospheric image motion displaces each position x_{im} by a small angle $\mathcal{M}_{im}$, which results in X_{im} being displaced by $\Delta\mathcal{M}_{im} = \sum_j a_{ij}(\mathcal{M}_{im} - \mathcal{M}_{jm})$. The spectral power density $G(q)$ of $\Delta\mathcal{M}_{im}$ in the space of spatial frequencies q is the product of two factors. The first factor $F'(q)$ depends on D , exposure time T , and properties of atmospheric turbulent layers generating image motion. The second factor depends only on the geometry of background star positions relative to the target i , and can be expanded in a series of even powers of q . Thus,

$$G(q) = F'(q) \sum_{s=1}^{\infty} q^{2s} F_{2s}(\tilde{x}_i, \tilde{y}_i; \tilde{x}_1, \tilde{y}_1 \dots \tilde{x}_N, \tilde{y}_N), \quad (5)$$

where $\tilde{x}, \tilde{y}$ are positions not affected by image motion (at zero turbulence). The coefficients F_{2s} are quadratic functions of coordinate differences that fastly decrease with s and take the form

$$F_{2s} = \sum_{\substack{\alpha, \beta=0 \\ \alpha+\beta=s}}^s A_{\alpha\beta}^{(2s)} \left[\sum_j a_{ij}(\tilde{x}_i - \tilde{x}_j)^\alpha (\tilde{y}_i - \tilde{y}_j)^\beta \right]^2 \quad (6)$$

where $A_{\alpha\beta}^{(2s)}$ are the characteristic constants. With the choice of coefficients a_{ij} , all sums in brackets can be turned to zero for a specific $A_{\alpha\beta}^{(2s)}$. In this way, the leading term F_2 and, possibly, several subsequent F_{2s} terms are eliminated, so that $G(k)$ will depend largely on the first non-zero component F_k of some optional mode $k = 4, 6 \dots$. For this purpose, the coefficients a_{ij} should satisfy each of $k(k+2)/8 - 1$ conditions

$$\sum_j a_{ij}(x_{i0} - x_{j0})^\alpha (y_{i0} - y_{j0})^\beta = 0, \quad \alpha + \beta = 1 \dots \frac{k}{2} - 1, \quad (7)$$

where $k \geq 4$, α and β are non-negative integers, and positions $\tilde{x}, \tilde{y}$ are substituted by respective measured positions of stars in the reference frame. The coefficients a_{ij} are determined for each target i by solving Eqs. (7) with normalizing condition (4), provided that at least $N' = k(k+2)/8$ reference stars are available. Because usually $N_m \gg N'$, the system of equations (7) is redundant and solvable with some useful restrictions on a_{ij} , for example Eq. (15). Due to the elimination of principal modes of the image motion spectrum, the resulting values of X_{im} are characterized by the residual image motion $\Delta\mathcal{M}_{im}$ that involves only uncompensated spectral modes of orders $s \geq k/2$ and therefore is much smaller than its initial value. Approximately, the variance of residual image motion is expressed by Eq. (1), where R is to be substituted by the radius R_i of local reference field for i -th target. The advantage of using larger orders in k is reflected in the coefficient B_k which decreases with k . In particular, for $k = 12$, its value is only 20% of that measured at the VLT with $k = 4$ (Paper I).

Next, let us introduce polynomial base functions $f_{1l} = 1$, $f_{2l} = x_{l0}$, $f_{3l} = y_{l0}$, $\dots$ $f_{wl} = x_{l0}^\alpha y_{l0}^\beta$ of coordinates x_{l0}, y_{l0} referring to the l -th star in the reference frame, with indices $w = 1, 2 \dots N'$ equal to the sequential number of combinations of α and β in equations (7). Taking advantage of the equality $\sum_j a_{ij}(x_{i0} - x_{j0})^\alpha (y_{i0} - y_{j0})^\beta = x_{i0}^\alpha y_{i0}^\beta - \sum_j a_{ij} x_{j0}^\alpha y_{j0}^\beta$, we can rewrite equations (4) and (7) in terms of functions f_{wl} as

$$\sum_j a_{ij} f_{1j} = 1, \quad \sum_j a_{ij} f_{wj} = f_{wi}, \quad w = 2, 3 \dots N' \quad (8)$$

3.2. Conversion from X, Y to displacements λ^x, λ^y

The image displacement X_{im} of the i -th star between frame m and the reference frame, filtered from low-order image motion and systematic distortions of the field, is measured relative to a set Ω_{im} of background stars whose positions are assumed to be fixed. However, since such are not available, we bypass this limitation by introduction of astrometric displacements λ that are related to all stars in the frame without distinction between reference and target objects. We consider X_{im} and λ related by

$$x_{lm} = x_{l0} + P_{lm} + \lambda_{lm}, \quad (9)$$

associating the measured position of stars in frame m with their position in the reference frame. Here, $P_{lm} = \sum_w c_{wm} f_{wl}$ is a polynomial function that contains powers of x, y not higher than $k/2 - 1$ and measured at discrete set of points x_{l0}, y_{l0} . Via coefficients c_{wm} , this function describes the systematic variation of stellar positions with the change of optical aberration and other similar effects. The second quantity

$$\lambda_{lm} = \hat{\xi}_l + e_{lm} - e_{l0} + h_{lm} - h_{l0} + \mathcal{M}_{lm} - \mathcal{M}_{l0} \quad (10)$$

is orthogonal to P_{lm} and satisfies conditions

$$\sum_i \lambda_{im}^x f_{wi} = 0, \quad \sum_i \lambda_{im}^y f_{wi} = 0, \quad w = 1, 2 \dots N' \quad (11)$$

In fact, Eqs. (9) and (11) give an expansion of x_{lm} into power series of f_{wl} , with λ_{lm} representing the remainders. λ_{lm} includes the “science” astrometric signal $\hat{\xi}_l$ (sum of proper motion, parallax, probable microlensing and reflex motion caused by the planetary companion) measured relative to the reference frame, random errors of photocenter measurement e_{lm} , the atmospheric chromatic displacement h_{lm} corrected by LADC, and the atmospheric image motion M_{lm} . These quantities (or their combinations) are also orthogonal to f_{wl} , with their low-order $w \leq N'$ expansion terms included in P_{lm} via coefficients c_{wm} .

The components of h_{lm} along the x - (opposite to right ascension) and y - (along declination) axes are given by (Paper I)

$$\begin{aligned} h_{lm}^x &= \rho_l \tan z_m \sin \gamma_m + d_l \tan \zeta_m \sin \gamma_m & \text{for } x\text{-axis} \\ h_{lm}^y &= -\rho_l \tan z_m \cos \gamma_m - d_l \tan \zeta_m \cos \gamma_m & \text{for } y\text{-axis} \end{aligned} \quad (12)$$

Here, ρ_l is the coefficient of differential chromatic displacement for the l -th star, d_l is a negative displacement produced by LADC to compensate for DCR effect, approximately of the same absolute size but opposite sign, z_m is a zenith distance at the midpoint of exposure of the m -th frame, γ_m is the parallactic angle, and ζ_m is the zenith angle of the LADC setting, which remains the same for a single series of observations.

Substitution of Eqs. (9) for x_{lm} and x_{jm} into Eq. (3) leads to elimination of the polynomials P_{lm} , while fulfilling the conditions given by Eq. (8). Taking into account that $X_{i0} = \sum_j' a_{ij}(x_{i0} - x_{j0}) = 0$ due to Eqs. (7), we obtain

$$\lambda_{im}^x - \sum_j' a_{ij} \lambda_{jm}^x = X_{im}. \quad (13)$$

An analogous expression for transformation of Y_{im} to λ_{im}^y results for measurements along the y -axis. Eqs. (13), for each frame m , form a system of $i = 1 \dots N_m$ linear equations with N_m unknowns λ_{im}^x . The solution for a given frame m is found irrespective of solutions for other frames. The matrix of the system (13), however, turns out to be degenerated due to N' linear dependences (7) between its columns (coefficients a_{ij}), and consequently it can be solved only using the same number of auxiliary restrictions, for example (11).

Below, we discuss two approaches to the extraction of $\hat{\xi}_i^x$, $\hat{\xi}_i^y$, ρ_i , and d_i from λ_{im} .

3.3. A case of fixed local reference star positions

In the case of small background star displacements λ_{jm} , and due to the averaging effect at summation, a term $\sum_j' a_{ij} \lambda_{jm}$ in Eq. (13) can be discarded as negligibly small. In fact, this assumes a fixed position of local reference stars $\lambda_{jm} = 0$ that leads to the most simple result $\lambda_{im}^x = X_{im}$. For each target i , we are facing a system of $2M$ equations

$$\begin{aligned} \xi_i^x + h_{im}^x - h_{i0}^x &= X_{im} - e_{im}^x - \Delta M_{im}^x \\ \xi_i^y + h_{im}^y - h_{i0}^y &= Y_{im} - e_{im}^y - \Delta M_{im}^y \\ m &= 1, 2 \dots M, \end{aligned} \quad (14)$$

for determining $\xi_i^x = \hat{\xi}_i^x - e_{i0}^x - \Delta M_{i0}^x$, $\xi_i^y = \hat{\xi}_i^y - e_{i0}^y - \Delta M_{i0}^y$, ρ_i , and d_i . This system can be solved by the least-squares method for each target object independently, with reference to its own subset Ω_{im} of reference stars, considering e_{im} and ΔM_{im} as random (with respect to index m) errors with variances σ_{im}^2 and σ_{at}^2 , respectively. Errors e_{jm} of reference star photocenter determinations add a further uncertainty to the phase of computation of X_{im} by means of Eq. (3). A variance of this error $\sigma_{it}^2 = \sum_j' a_{ij}^2 \sigma_{jm}^2$,

equal to the variance of a term $\sum_j' a_{ij} x_{jm}$ in Eq. (3), can be reduced applying restrictions

$$\sum_j' a_{ij}^2 \sigma_{jm}^2 = \min \quad (15)$$

on solutions a_{ij} of the system (7).

As it was mentioned above, geometric field distortions (their variations in time), up to order k , are excluded along with image motion.

This simple method was applied for the test reduction of four-hour series of frames obtained at VLT/FORS2 during one night (Paper I). By filtration of atmospheric image motion carried out with k up to 12, 300 μs precision of a single position for bright stars was achieved, which is near to the accuracy of photocentre measurements. Approximation $\lambda_{jm} = 0$ was valid due to a small residual DCR effect for frames obtained with the R_{special} filter.




Fig. 1. x , y residuals of Eqs. (14) for 25 images computed with the assumption of “fixed” background star positions (*left panel*) in comparison to the exact solution (*right*) for epochs of reference and processed frames being separated by two years. Dot sizes are proportional to the stellar magnitude (only bright $B < 21.5$ mag stars are shown). Note the difference in scales (1 mpx = 100 μs).

3.4. Reduction for floating reference stars

For the processing of sky images in the B -filter, the use of the technique described above is however limited, even for frames taken within the same night. In this case, an excess of positional errors over σ_i by a factor 3–5 results, because the underlying assumption of $\lambda_{jm} = 0$ does not hold due to large (~ 2 –10 mas/hour) chromatic displacement of images. For frames taken two years after the reference frame, which results in significant additional proper motion displacements, the solution of Eqs. (14) produces unacceptably large residuals (Fig. 1a).

The exact solution of Eqs. (11–13) ensures a linearity and uniqueness of the transformation from X_{im} to λ_{im} even for moving ($\lambda_{im} \neq 0$) reference stars. Therefore, although X_{im} are determined relative to local reference fields Ω_{im} centered at the i -th target and depend on the size R_i of these fields, the final solutions λ_{im} are consistent within the global reference field of size R and do not depend on the choice of initial R_i values. Since usually $R > R_i$, the transformation from X_{im} to λ_{im} increases the atmospheric image motion component in λ_{im} , which should be compensated for by moving to larger k .

Note that due to conditions (8), both λ_{im} and $\lambda_{im} + \varphi_{im}$ are the solutions of a system (13) if φ_{im} are functions

$$\varphi_{im}^x = \sum_{w=1}^{N'} E_{wm}^x f_{wi}, \quad \varphi_{im}^y = \sum_{w=1}^{N'} E_{wm}^y f_{wi} \quad (16)$$

with arbitrary coefficients E_{wm}^x and E_{wm}^y . Therefore, for splitting λ_{im} into components we can use

$$\begin{aligned} \xi_i^x + h_{im}^x - h_{i0}^x + \varphi_{im}^x &= \lambda_{im}^x + \mathcal{N}_{im}^x \\ \xi_i^y + h_{im}^y - h_{i0}^y + \varphi_{im}^y &= \lambda_{im}^y + \mathcal{N}_{im}^y \\ m = 1, 2 \dots M, \quad i = 1, 2 \dots N \end{aligned} \quad (17)$$

where $\mathcal{N}_{im} = -e_{im} - \Delta M_{im}$ is a noise with variance $\varepsilon_{im}^2 + \sigma_{at}^2$. Eqs. (17) can then be solved as a system of $2 \sum N_m$ conditional equations with $4N + 2MN'$ unknowns ξ_i^x , ξ_i^y , ρ_i , d_i , E_{wm}^x , and E_{wm}^y .

Because for most not very faint targets, $\sigma_{at} \ll \varepsilon_{im}$, the least-squares solution is found with weights $g_{im} = \varepsilon_{im}^{-2}$. Also, due to the dominant contribution of bright stars, we can safely assume $g_{im}^2 \approx I_i / \theta_{im}^2$ which follows from the approximation given by Eq. (2) for these stars.

Linear relations between some parameters of a system (17) do not permit to separate them completely. For example, systematic variations of ξ_i^x , ρ_i , and d_i quantities over a field imitate the behaviour of φ_{im}^x functions. Associations of this type are eliminated by applying N' orthogonalizing restrictions

$$\begin{aligned} \sum_i \xi_i^x f_{wi} I_i &= 0, \quad \sum_i \xi_i^y f_{wi} I_i = 0, \\ \sum_i \rho_i f_{wi} I_i &= 0, \quad \sum_i d_i f_{wi} I_i = 0, \quad w = 1, 2 \dots N' \end{aligned} \quad (18)$$

The restrictions (18) permit us to recover only those components of ρ_i , d_i , ξ_i^x , and ξ_i^y , that do not correlate with functions f_{wi} .

An essential drawback of a straightforward solution of Eqs. (16-18) is a large number of unknowns E_{wm}^x and E_{wm}^y , that cause a certain instability of a solution. For example, the number of unknowns for Dec 2000 frames is $2MN' = 2 \cdot 40 \cdot 36 \approx 2900$ at $k = 16$, which is comparable to the number of measurements $2 \sum N_m \approx 9100$.

However, there is a simpler solution. Consider equation with a diagonal element E_{wm}^x of the normal system of Eqs. (17):

$$\begin{aligned} \sum_{i \in \omega_m} (\xi_i^x + h_{im}^x - h_{i0}^x) f_{wi} I_i + \sum_{i \in \omega_m} f_{wi} I_i \sum_w E_{wm}^x f_{wi} \\ = \sum_{i \in \omega_m} \lambda_{im}^x f_{wi} I_i \end{aligned} \quad (19)$$

where summation is performed only over a limited sample ω_m of stars measured in the m -th frame and, since $\text{FWHM} \approx \text{const}$ within the frame, the approximation $g_{im} = I_i$ was used. Adding the sum

$$\Lambda_{wm}^x = \sum_{i \in \text{not } \omega_m} (\xi_i^x + h_{im}^x - h_{i0}^x) f_{wi} I_i \quad (20)$$

formed over stars *not* measured in the frame m to both sides of Eq. (19), and taking into account (18), we note that the first component in Eq. (19) turns to zero. Therefore

$$\sum_{i \in \omega_m} f_{wi} I_i \sum_w E_{wm}^x f_{wi} = \sum_{i \in \omega_m} \lambda_{im}^x f_{wi} I_i + \Lambda_{wm}^x. \quad (21)$$

If a solution of the system (13) for the m -th frame is found that fulfills the conditions

$$\sum_{i \in \omega_m} \lambda_{im}^x f_{wi} I_i = -\Lambda_{wm}^x \quad w = 1, 2 \dots N', \quad (22)$$

instead of those of Eq. (11), the right part of the system given by Eq. (21) turns to zero, thus yielding a unique solution $E_{wm} = 0$. Similar considerations for the y -component of the data yield $E_{wm} = 0$, hence $\varphi_{im}^x = \varphi_{im}^y = 0$. The elimination of the large number ($2MN'$) of unknowns E_{wm}^x and E_{wm}^y in Eq. (17) occurs due to imposing an equivalent number of restrictions (22), thereby increasing the effective degree of freedom of the system given by Eq. (17) to $2 \sum N_m - 4N$. Furthermore, this splits the system given by Eq. (17) into N independent subsystems of $2n_i$ equations (n_i is number of observations of the i -th star)

$$\begin{aligned} \xi_i^x + h_{im}^x - h_{i0}^x &= \lambda_{im}^x + \mathcal{N}_{im}^x \\ \xi_i^y + h_{im}^y - h_{i0}^y &= \lambda_{im}^y + \mathcal{N}_{im}^y \end{aligned} \quad (23)$$

which is easily solved for ξ_i^x , ξ_i^y , ρ_i , and d_i for each of the star observed independently. With this processing technique, Λ_{wm} entering Eq. (22) are calculated by iteratively refining ξ_i^x , ξ_i^y , ρ_i , and d_i . Iterations are however not required if each star is measured in all of the frames since in this case the identity $\Lambda_{wm}^x = \Lambda_{wm}^y = 0$ holds due to Eq. (18).

Some comments should be made on the nature of ξ values computed not in the absolute reference frame, but only relative to a limited set of N nearby stars. Systematic differences between ξ and values ξ_{abs} referring to the absolute or global frame, are given by

$$\xi_{abs,i} = \xi_i + \sum_w G_w f_{wi} \quad (24)$$

which incorporates a polynomial function $\sum_w G_w f_{wi}$ of N' order with coefficients G_w . The values G_w should confirm the restrictions given by Eq. (18) which now take a form $\sum_i (\xi_{abs,i} - \sum_w G_w f_{wi})^2 I_i = \min$, or $\sum_i \xi_i^2 I_i = \min$. Eq. (24) thus is an expansion of $\xi_{abs,i}$ into base functions f_{wi} , and ξ_i are the residuals of the expansion. The increase of N' (or k) leads to the increase of polynomial modes excluded from ξ_{abs} and thus to the loss of information retained in ξ . Reduction therefore should use the lowest k that still ensures acceptably small σ_{at} . Above considerations, of course, are fully applicable to ρ_i and d_i parameters as to the remainders of corresponding polynomial expansions.

4. Reduction of measured images

An image obtained on 21 Dec 2000 at an hour angle near to the mean hour angle of other images was taken as a reference frame. Only stars in the central region of the sky with an $R = 1000$ px radius containing $N = 169$ star images down to $B = 24$ mag were selected for the reduction. Peripheral stars which due to a strong jittering were often beyond frame boundaries thus were rejected. Nevertheless, the number of stars measured varied among the frames and reached down to $N_m \approx 90$.

We have taken advantage of the facts that the colours of the observed stars do not vary significantly on short time-scales (few days) and positions of most stars are only slightly affected by proper motions and parallaxes. Therefore frames were combined into series within annual epochs, which statistically improved the accuracy of model parameters determination. We therefore organized our data into two series of 39 and 25 frames for 2000 and 2002-2003 epochs, respectively, each one representing four nights. Astrometric reduction was carried out with different choices of the parameter k , where the accuracy of results became sufficiently good at $k = 16$, which was adopted for the final reduction. Our iterative procedure involved the computation of ξ_i^x , ξ_i^y , ρ_i , as well as d_i and the refinement of Λ_{im} quantities calculated on the basis of the former parameters. The number

of outliers at each coordinate was limited to 1.25% of the total number of measured star images.

The residuals u_{im}^x and u_{im}^y of the conditional equations Eq. (23) on x - and y - coordinates are uncorrelated (Fig.1b) and, besides, substantially smaller than those obtained with the approximation of "fixed" background star positions (Fig.1a). We also did not find any correlation between u_{im} and the stellar position within the frame or the colour-dependent parameters ρ_i and d_i . As a simple illustration for a further discussion, Fig.2 presents examples of u_{im}^x residuals for stars of different magnitude, from bright $B = 18.5$ to faint $B = 23$ mag. A difference in the data point scattering reflects the dependence of photocenter error on the light signal. No apparent correlation in time is seen. More extensive statistical analysis of u_{im} values is performed in Sect.5.




Fig. 2. Example of u_{im}^x residuals for 2000 epoch frames.

4.1. Performance of the LADC

Table 2. Relative corrections to $\tan \zeta_m$: producing best accuracy of observations (C_m) and corresponding to a 10 mn delay in the LADC setting (C'_m)

night	C_m	C'_m	night	C_m	C'_m
2000			2002-2003		
20 Dec	-0.17	-0.15	29 Dec	-0.06	-0.11
21 Dec	0	-0.02	30 Dec	0.05	0.09
22 Dec	0.26	0.20	31 Dec	-0.11	-0.15
23 Dec	-0.11	-0.14	02 Jan	0.02	-0.13

Zenith distances ζ_m of the LADC setting used in Eqs. (12) are not specified in the fits file headers. According to the technical documentation of the VLT, ζ_m is equal to the telescope zenith distance at the midpoint exposure time of the first science image of the series and is constant for all frames within the series. We however tried to use corrected values $(1 + C_m) \tan \zeta_m$ instead of $\tan \zeta_m$ in order to improve internal precision. Coefficients C_m (Table 2) that yield the best-achievable precision have been determined with an accuracy of about ± 0.03 relative to some zero-point night, chosen as 21 Dec 2000. Closer examination revealed that $(1 + C_m) \tan \zeta_m$ is approximately equal to $\tan \zeta'_m$ computed with the telescope zenith distance ζ'_m 10 mn prior to the first exposure. Writing ζ'_m in the form $\tan \zeta'_m = (1 + C'_m) \tan \zeta_m$ allowed us to introduce new values C'_m and compute them in analogy to

C_m . A good consistency of C_m and C'_m values suggests that actual angles ζ_m are really close to ζ'_m . According to Jehin E. (private communication, 2006), this may occur when the LADC is not reset after finishing preceding observations of the same target. The final reduction was carried out with $(1 + C_m) \tan \zeta_m$ rather than $\tan \zeta_m$.




Fig. 3. The LADC compensating displacement d_i and atmospheric differential chromatic refraction ρ_i for stars of $B = 18 - 24$ mag (dots) with a linear (solid line) fit. The dot size is proportional to the stellar magnitude. Individual error bars are shown.

An excellent performance of the LADC is demonstrated by Fig.3 which reproduces the distribution of ρ_i and d_i (averaged over both epochs) with error bars based on the effective precision of a single measurement (26). Blue stars are shown in the upper left corner. The distribution of stars is strongly concentrated along a line $\rho_i = -1.2d_i$ with a scatter that only slightly exceeds random errors.

4.2. Effective precision of a single measurement

Astrometric positions are affected by various sources of error. The most important are fundamental (photon noise in the image), of instrumental origin (optical field distortions, errors related to CCD, etc.), induced by the atmosphere (DCR effect, image motion), or related to inefficient reduction techniques that should take into account and exclude largest errors. Without analyzing each source separately, some of which are rather efficiently filtered out at the phase of the reduction, we consider their total impact on the key system of Eqs. (23), whose solution yields final astrometric parameters ξ_x and ξ_y . This total error in Eqs. (23) was denoted as $\mathcal{N}_{im}$ and includes e_{im} , $\Delta \mathcal{M}_{im}$, and a sum of other residual errors not filtered by the reduction model. $\mathcal{N}_{im}$ includes both a random noise and components correlated in time. The presence of correlated components leads to strongly biased results if the correlation time exceeds the length of a single series, while being shorter than duration of the observational campaign. In this case, processing a short series of frames will recover noise signature rather than the desired real astrometric signal. In next section we described results of a statistical analysis of $\mathcal{N}_{im}$ based on measured residuals u_{im} .

The variance σ_{im}^2 of the noise $\mathcal{N}_{im}$ is the effective variance of a single measurement. Obviously, $\sigma_{im}^2 \geq \varepsilon_{im}^2 + \sigma_{at}^2$. The size of σ_{im} depends on the stellar magnitude I_i , the seeing θ_{im} , the sky




Fig. 4. Effective precision of a single measurement $\bar{\sigma}$ (mean for x and y axes) for 2000 (circles) and 2002-2003 (triangles) epochs as a function of B . Centroiding error $\bar{\varepsilon}$ at average seeing and background noise is shown by solid (first epoch) and dashed (second epoch) lines. A scale is $1 \text{ mpx} = 100 \mu\text{as}$.

level b_m , the reduction parameters k and R , etc. In the following, the analysis simplifies by using the improved expression for ε_{im} that, as compared to Eq. (2), more adequately takes into account variations of θ_{im} and b_m , amounting to about $\pm 30\%$, and is valid in a wide (about 5 mag) range of stellar brightness. Considering asymptotic dependences of ε on θ and b given by Irwin (1985) for bright and faint images, we modified Eq. (2) to

$$\varepsilon_{im} = \frac{\theta_{im}}{2.46 \sqrt{I_i}} \left(1 + \alpha_0 I_i^{-0.7} \frac{\theta_{im}}{\theta_0} \sqrt{b_m/b_0} \right). \quad (25)$$

The second component in parentheses is significant for faint $B > 21.5$ stars only. The θ_{im} , or FWHM value, is available as a PSF model parameter (Sect.2). We have found that the variance of the normalized deviations u_{im}/ε_{im} based on our new relation is almost independent from observation conditions. For that reason, the actual observed value of σ_{im}^2 was obtained as a systematic correction to ε_{im}^2 :

$$\sigma_{im}^2 = \varepsilon_{im}^2 \left\langle \frac{1}{n_i - 2} \sum_m u_{im}^2 / \varepsilon_{im}^2 \right\rangle \quad (26)$$

where an average was taken over stars of approximately the same magnitude. Further averaging of σ_{im}^2 with respect to i and m (over observations) yielded the effective variance of a single measurement $\bar{\sigma}^2$ at average observational conditions during the considered epoch. Fig.4 presents both $\bar{\sigma}$ and the mean centroiding error $\bar{\varepsilon}$ corresponding to average observing conditions as a function of B . Because $\bar{\sigma}$ matches well the errors $\bar{\varepsilon}$ predicted by our numerical simulation, we conclude that the sum of all errors arising from different sources, including σ_{at} , is small in comparison to ε , so that, approximately, $\bar{\sigma} \approx \bar{\varepsilon}$. This means that the astrometric precision depends only on errors of the photocentre determination which are random Gaussian and uncorrelated with time due their nature of arising from photon statistics. The latter property is very important since it forms the basis for substantial statistical improvements of the astrometric accuracy $\Delta(n_i)$ for a sequence of n_i frames with the increase of the number of measurements as

$$\Delta(n_i) = \bar{\varepsilon} / \sqrt{n_i}; \quad n_i \leq n_{\max} \quad (27)$$

where $\bar{\varepsilon}$ is the photocenter error at average observation condition. The estimate (27) is approximate since assumes the best situation with a diagonal covariance matrix of the system (23). The validity of Eq. (27) is limited with respect to the maximum amount of frames $n_{\max}$, at which the value of $\Delta(n_i)$ reaches a floor set by systematic errors.

5. Testing the VLT temporal astrometric stability

5.1. The Allan variance

The simplest empiric validation of Eq. (27) and illustration of how this law works at different n_i , is associated with use of Allan variance σ_{Al}^2 of the residuals u_{im} . This quantity (Pravdo & Shaklan 1996) is a function of time lag $\tau \leq n_i/2$ expressed in frames

$$\sigma_{Al}^2(\tau) = \frac{1}{2(n_i + 1 - 2\tau)} \sum_{p=1}^{n_i+1-2\tau} \left[\tau^{-1} \sum_{q=0}^{\tau-1} (u_{p+q} - u_{p+q+\tau}) \right]^2 \quad (28)$$

$\sigma_{Al}^2(\tau)$ is a precision of the average of u_{im} taken over τ frames, and so corresponds to $\Delta^2(\tau)$. For uncorrelated sets of u_{im} , the expected dependence $\sigma_{Al}^2 \sim \tau^{-1}$ is consistent with Eq. (27). In effort to increase the range of time lags, we performed computations of σ_{Al}^2 treating all epoch frames as a single series of n_i length. For miscellaneous reasons, n_i varied from about $0.4M$ to $1.0M$. Results for stars of different B magnitude classes are shown in Fig.5 (two bottom graphs). Though, in general, plots for separate stars follow the expected law τ^{-1} (straight lines with an ordinate $\bar{\varepsilon}^2$ at $\tau = 1$), it is difficult to make any conclusion at $\tau \geq 10$. To smooth statistical fluctuations and exclude the dependence on the star brightness, we normalized individual σ_{Al}^2 by $\bar{\varepsilon}^2$ (upper plot of Fig.5) and then performed averaging of results over all 169 stars. Obtained smooth functions $\langle \sigma_{Al}^2 / \bar{\varepsilon}^2 \rangle$ (black dots) indicate no evidence of large systematic error, and extend a validity of the approximation $\sigma_{Al}^2 \sim \tau^{-1}$ at least to $\tau \approx 15$. Hence Eq.(27) is valid to about $n_{\max} \approx 15$. The asymptotic floor of the Allan variance at large τ is under the detection limit of this approach. We can claim however that with 15 frames, the resulting precision of the average is about $50 \mu\text{as}$ for bright stars. In Sect.5.3, we renew this discussion to derive more exact estimate of $n_{\max}$.

5.2. Distribution of residuals

The null hypothesis H_0 formally arising from above analysis is that the $\mathcal{N}_{im}$ component in Eqs. (23) represents an uncorrelated Gaussian noise. The ensuing consequences of this assumption are

- the averaging law, Eq. (27), and
- the absence of other noise components, including instrumental errors, with a noticeable magnitude.

We applied several statistical tests of our hypothesis. In the first test, we formed a histogrammic distribution W of normalized deviations $u'_{im} = \sqrt{n_i/(n_i - 2)} u_{im} / \sigma_{im}$ and compared it with the distribution W_0 of these residuals expected when the data fulfill the hypothesis H_0 . In order to obtain W_0 , we performed some numerical simulations of observations with a typical spacing of frames on hour angles, and adding a Gaussian noise with a unit variance in x and y to the right side of Eqs. (23). Observed distributions W were computed using all combinations of data sets available (2000 and 2003/2003 epoch frames, x and




Fig. 5. Allan variance as a function of time lag τ for individual stars (dashed lines) in comparison to the τ^{-1} law (straight lines) in narrow magnitude ranges (two lower graphs) and the normalized variance $\sigma_{AI}^2/\varepsilon^2$ for all stars (upper plot). The average $\langle \sigma_{AI}^2/\varepsilon^2 \rangle$ taken over all stars is shown by black dots. The plots refer to x coordinate and 2000 epoch frames.

y data, with faint, bright, or a whole sample of stars). Fig. 6a, b present an example of such a histogram W for bright stars. Corresponding simulated distributions W_0 are shown by smooth curves, for which the same binning as for the computation of W has been applied. Unlike N_{im} , the distributions W_0 and W are not Gaussians and have a more strongly peaked shape. The consistency of observed and model distributions by visual inspection is rather good. Numerically, it was estimated by χ^2 values computed from differences between W and W_0 in fixed binned intervals containing at least 5–10 data points. The χ^2 values with corresponding degrees of freedom (Dof) are listed in Table 3 for faint, bright, or all stars, respectively. The χ^2 -test based on these data does not reveal a deviation of W from W_0 at significance levels below 20%, which does not allow to reject our hypothesis H_0 . Due to a limited number of data points, this test is however only sensitive to a central $\pm 2\sigma$ region of the distribution.

Table 3. χ^2 values and Dof characterizing differences between W and W_0

	faint	bright	all stars
2000 epoch			
$\chi^2, x/y$	20.1/21.5	23.0/22.7	24.4/28.6
Dof	22	22	24
2002/2003 epoch			
$\chi^2, x/y$	20.8/11.6	25.5/17.2	27.8/27.1
Dof	18	20	22




Fig. 6. Histograms of individual normalized residuals u'_{im} in x (a) and in y (b) for actual (steps) and simulated (smooth lines) observations, on 2000 (solid) and 2002 epochs (dashed lines); the same for normalized night average residuals $\bar{u}_{ip}/\sigma_p$ (c and d). Plots are drawn for bright $B < 20.5$ stars.




Fig. 7. The differences D_n between the observed and model cumulative frequency distributions of normalized residuals u'_{im} for 2000 epoch measurements in y (bright stars)

We applied the more powerful Kolmogorov-Smirnov test to detect a potential difference between the observed and simulated distributions. For this purpose, we computed cumulative frequency distributions of observed and model normalized residuals u'_{im} , and then obtained the difference D_n between two distributions. A case with largest deviations for 2000 epoch measurements in y with 1721 data points (bright stars only) is shown in Fig. 7 where the differences D_n are plotted as a function of the

normalized deviation. The Kolmogorov-Smirnov test based on this plot (in this worst case $\max|D_n| = 0.024$) shows no significant difference between the model and measured distributions at 20% confidence level. Both tests applied thus do not reject hypothesis H_0 of a really normal distribution of N_{im} at approximately the same significance levels.

For another test, we considered nightly averages $\bar{u}_{ip} = \sum_{m \in p} u_{im} \sigma_{im}^{-2} / \sum_{m \in p} \sigma_{im}^{-2}$, involving all observations of the i -th star within a single p -th night. Fig.6 c, d show observed and model frequency distribution of normalized residuals $\bar{u}_{ip}/\sigma_p$, where σ_p is the error of the determination of $\bar{u}_{ip}$. A good agreement of the histograms of the observed statistics (both for bright and faint stars) with their expectations from model simulations, a symmetry, and the absence of large deviations (normally less than 1% of $\bar{u}_{ip}/\sigma_p$ values deviate at about 3 units) demonstrate excellent stability of the VLT instrumental system during time spans of a few nights. Compatibility of the observed and model distributions is also confirmed by the Kolmogorov-Smirnov- and χ^2 -tests.

5.3. Integral statistics

Table 4. Integral statistics S_0 and S_A for bright $B < 20.5$, faint $B > 20.5$ stars, and for simulated observations

	S_0^x	S_0^y	error	S_A^x	S_A^y	error
2000 epoch						
model	0.997	0.985	-	0.749	0.839	-
bright	0.992	0.989	± 0.021	0.784	0.849	± 0.064
faint	0.982	0.988	± 0.019	0.761	0.787	± 0.054
2002-2003 epoch						
model	0.994	0.973	-	0.738	0.845	-
bright	0.973	0.961	± 0.029	0.722	0.872	± 0.067
faint	0.989	0.952	± 0.026	0.744	0.874	± 0.055

We also considered integral statistics $S_0^2 = \langle \frac{1}{n_i - 4} \sum_m \frac{(u_{im} - \bar{u}_{ip})^2}{\sigma_{im}^2} \rangle$, where averages are taken over all stars and nights, and $S_A^2 = \langle \frac{1}{4} \sum_p \bar{u}_{ip}^2 / \sigma_p^2 \rangle$, where an average is taken over all stars. These two statistics are either sensitive to the variation of systematic errors within a given night (S_0) or from one night to another (S_A), and therefore are a powerful indicator of their presence. The quantities S_0 and S_A , calculated separately for both coordinates x and y , are listed in Table 4 for each epoch of observations together with the respective expected values $S_0(\text{mod})$ and $S_A(\text{mod})$, obtained from numerical simulations where random Gaussian noise N_{im} has been assumed. Errors due to a limited number of measurements indicate an 80% confidence interval for possible deviations of observed values from our model assumption. Observed statistics match well the model within the error limits.

The fact that we do not find an excess in the measured S_0 and S_A values means that no or very small extra noise except N_{im} could be present in measurements. It is useful to estimate its upper limit. Suppose that the observed positions, besides of N_{im} , are affected by some small systematic error that is constant within a night but varies between nights with an amplitude peculiar to a certain star. The observed value of $S_A^2(\text{obs})$ then exceeds its model expectation $S_A^2(\text{mod})$ by a small amount S_{sys}^2 which is the average signal variance for a given star sample. These quan-

ties are related by $S_A^2(\text{obs}) = (\chi_{\text{Dof}}^2 / \text{Dof}) S_A^2(\text{mod}) + S_{\text{sys}}^2$. For statistical reasons, it is probable that the sampled value $S_A^2(\text{obs})$ is equal or below of its mathematical expectation $S_A^2(\text{mod}) + S_{\text{sys}}^2$. The largest value of S_{sys}^2 , according to Table 4, can exist in the x -axis measurements of bright stars (2000 epoch), for which we found $S_{\text{sys}}^2 / S_A^2(\text{mod}) \leq 0.19$ at 20% confidence level. This is a ratio of a systematic to random error component for a single night series represented, in average, by 8 frames. In other words, the bias related to systematic errors is equal to a random component expected in positions averaged over $8/0.19 = 40$ frames. This is an estimate of n_{max} introduced in Eq.(27), more precise, as compared to that obtained in Sect.5.1. For y -coordinate, we obtain $n_{\text{max}} = 65$. Frames of 2002 yield $n_{\text{max}} = 120$ and 50 for the x and y axes correspondingly. With $n_{\text{max}} = 50$ adopted as a reliable limit, precision of a night series is about $30 \mu\text{s}$.

5.4. Correlations

A final important test was carried out to investigate correlations in N_{im} which can negatively affect the averaging law (27). For that purpose, we computed the autocorrelation function of normalized residuals

$$r(\tau) = H(\tau) \langle (u_{im_1} / \sigma_{im_1}) (u_{im_2} / \sigma_{im_2}) \rangle \quad (29)$$

of the argument $\tau = m_2 - m_1$, which is equal to the difference of frame indexes and used as a time lag. Here, $H(\tau) = 1 - \tau/M$ is the weight function that compensates the increase of statistical variations at large τ . Averaging is performed over the images of all stars. A corresponding computation of the expected correlation function $\hat{r}(\tau)$ was carried out with the model observations containing white noise with a unit variance. The observed $r(\tau)$ values computed for both coordinates x and y for bright $B < 20.5$ mag stars are shown in Fig.8 along with the model function $\hat{r}(\tau)$ (mean for x and y) and $1-\sigma$ limits for statistical scatter of individual deviations. Oscillations in $\hat{r}(\tau)$ plot inherit a compound four-night structure of the series. Observed data in general follow the expected dependence, though with few isolated deviations over 2 sigma.




Fig. 8. Observed autocorrelation function in x (circles) and in y (squares) as compared to the model function (solid line) with a $1-\sigma$ tolerance limits (two dotted lines) and approximation (30) (dashed line)

Both r and $\hat{r}$ are not identically zero as it could be expected for uncorrelated measurements. The observed systematic negative bias of correlations, however, is not due to instrumental errors but has a theoretical background (Lazorenko 1997). In order to see this, consider Eqs. (23) for the i -th star. When the

number of frames M is large and the distribution of frames with hour angle is sufficiently random, all parameters in the system given by Eq. (23) become uncorrelated. Then ξ_i^x is approximately equal to the average of $(\lambda_{im}^x - h_{im}^x + h_{i0}^x + N_{im})$ taken with respect to the index m . In this approximation, one finds $u_{im}^x = (\lambda_{im}^x - h_{im}^x + h_{i0}^x + N_{im}) - \xi_i^x$. The term given in parentheses formally represents a solid time series of the "time" argument m with $m = 1 \dots M$ measurements containing noise. With this approach, the u_{im}^x values are the remainders of a time series after subtraction of the fitting polynomial (namely a constant ξ_i^x in our case). The bias of the autocorrelation function shape due to subtraction of the best-fitting polynomial for measurements limited in time was also previously studied by Lazorenko (1997). The autocorrelation function r' of series remainders takes a particularly simple form if the power spectrum of measurement errors resembles a white noise (which is the case) with a unit variance and constant rate of data sampling. In this case, one obtains (Bakhonski et al. 1997)

$$r'(\tau) = -\frac{2\zeta}{M} \text{sinc}(2\pi\zeta\tau/M) \quad (30)$$

where $\zeta = 0.56$, if the subtracted polynomial is presented by a constant, and $\text{sinc}(z) = \sin(z)/z$. One can see that $r' \approx 0$ for large M only (long series) while for small M (limited number of sampled data) its value is absolutely large and negative. Due to simplification of our considerations, the measured and model functions are not exactly fitted by r' , but good enough however to explain the statistical origin of the observed negative bias (Fig. 8). We conclude that no correlation is present in N_{im} at scales of several frames.

Table 5. Observed $\bar{r}$ and model $\bar{r}_{\text{mod}}$ normalized correlations.

τ'	$\bar{r}^x$	$\bar{r}_{\text{mod}}^x$	$\bar{r}^y$	$\bar{r}_{\text{mod}}^y$	1-sigma error
2000 epoch					
1	-0.17	-0.18	-0.22	-0.24	± 0.05
2	-0.17	-0.10	-0.20	-0.24	± 0.07
3	-0.42	-0.38	-0.25	-0.23	± 0.09
2002-2003 epoch					
1	-0.15	-0.14	-0.26	-0.25	± 0.06
2	-0.19	-0.20	-0.32	-0.23	± 0.07
3	-0.35	-0.26	-0.13	-0.19	± 0.10

Correlation at time scales of few days were studied using normalized nightly-average residuals $\bar{u}_{ip}/\sigma_p$. We computed correlations

$$\bar{r}(\tau') = \langle (\bar{u}_{ip_1}/\sigma_{p_1})(\bar{u}_{ip_2}/\sigma_{p_2}) \rangle \quad (31)$$

as a function of the time lag $\tau' = p_2 - p_1$ between two nights with indexes p_1 and p_2 , expressed in days. The values of $\bar{r}$ for the observed data and corresponding model value $\bar{r}_{\text{mod}}$ are given in Table 5. Apparently, there is a strong negative bias in $\bar{r}$ due to a very limited number of data points in our 4-night series. This bias, however, is well-modelled using uncorrelated noise N_{im} , which confirms previous conclusions that there are no traces of instrumental signature in VLT observations at time scales of few nights, and supports the validity of the averaging law (27).

5.5. Two-year stability

Given only two epoch measurements, no conclusions on a long-term astrometric stability of the VLT can be made based on




Fig. 9. Difference in the chromatic parameters ρ_i and d_i between the two epochs, normalized by the error of this difference computation (dots). Ellipses are drawn at 1% (dashed line) and 5% (solid) significance level. A straight line shows the direction of point displacements due to the actual change of star colours.

positional information. Considering extreme importance of this problem, we present here indirect analysis of this issue based, instead of positions, on a comparison of chromatic ρ_i and d_i parameters computed from independent runs of 2000 and 2002/2003 epochs. If one assumes constant colours for the observed stars, these parameters should coincide within error bars. A change of the VLT characteristics, depending on stellar colours, should result in an additional difference. Fig. 9 shows normalized differences between ρ_i and d_i between the two epochs, where ellipses refer to significance levels of 1% or 5%, respectively. Some points represent outliers, testifying the presence of systematic errors. The largest residuals for bright stars in the right bottom however occur along a line $\rho_i = -1.2d_i$ that is peculiar to the distribution of chromatic parameters shown in Fig. 3. It is therefore very likely that these deviations are caused by a change in stellar colour. Some other large deviations appear to be related to stars in outer regions of the observed field where the method seems to give biased results. Except for these outliers, the distribution of points in Fig. 9 resembles a Gaussian, which is an indicator of a rather good astrometric stability of the VLT even at very long time scales. It should be stressed that above test is insensitive to non-chromatic type of errors and therefore is only indicative.

6. Conclusion

Our results reveal an exceptionally good astrometric performance of the VLT and its camera FORS1. Thus, the precision $\bar{\varepsilon}$ of the position of a bright star reaches $200 \mu\text{as}$ for a single measurement, which fairly well corresponds to the former $300 \mu\text{as}$ estimate obtained for FORS2 camera (Paper I). Both cameras thus give similarly good results. The term "bright star" refers to unsaturated images containing $1\text{--}3 \cdot 10^6$ electrons and may correspond to different stellar magnitudes depending on the exposure, filter, and seeing. Most importantly, we found negligibly small systematic errors of instrumental and other origin. In fact, no traces of these 'dangerous' errors were found at time intervals up to 4 days. Due to this fact, the precision for a frame series improves as $\bar{\varepsilon}/\sqrt{n_i}$, and at $n_i = 50$ reaches $30 \mu\text{as}$ for FORS1 and

40 μ as for FORS2. With use of FORS2, the observation time needed to obtain this accuracy is about 1 hour if short 15 s exposures and a 2×2 binning of pixel reading is used. We conclude that the VLT with cameras FORS1/2, due to its enormous collecting light power, fine optical performance, and effective averaging of wave-front distortions over a large aperture, is a powerful instrument that can be used efficiently for high-precision astrometric observations of short-term events, in particular, of planetary microlensing.

Acknowledgements. We would like to thank Dr. E.Jehin for his helpful comments on the details of LADC operation.

References

- Avila, G., Rupprecht, G. & Beckers, J. M. 1997, *Proc. SPIE* 2871, 1135
 Bakhonski, A.V., Kostjutshenko, V.L., & Lazorenko, P.F. 1997, *Kinematics and Phys. Celest. Bodies* 13, 5, 50
 Beaulieu, J.-P., Bennett, D.P., Fouque, P., et al. 2006, *Nature*, 439, 437
 Boden, A.F., Shao, M., & Van Buren, D. 1998, *ApJ* 502, 538
 Bond, J.A., Udalski, A., Jaroszynski, M. et al. 2004, *ApJ* 606, L155
 Colavita, M.M., Boden, A.F., & Crawford, S.L. 1998, *Proc. SPIE* 3350, 776
 Dominik, M. 1999, *A&A* 349, 108
 Dominik, M., & Sahu, K.C. 2000, *ApJ* 534, 231
 Irwin, M.J. 1985, *MNRAS*, 214, 575
 Jaroszyński, M., & Paczyński, B. 2002, *Acta Astron.* 52, 361
 Gaudi, B.S., & Gould, A. 1997, *ApJ* 486, 85
 Gaudi, B.S. 1998, *ApJ* 506, 533
 Gaudi, B.S., & Han, C. 2004, *ApJ* 611, 528
 Gould, A., Udalski, A., Bennett, D., et al. 2006, *ApJ* 644, L37
 Han, C., & Chunguk, L. 2002, *MNRAS*, 329, 163
 Hardy, S.J., & Walker, M.A. 1995, *MNRAS* 276, L79
 Høg, E., Novikov, I.D., & Polnarev, A.G. 1995, *A&A* 294, 287
 Lane, B.F., & Muterspagh, M.W. 2004, *ApJ*, 601, 1129
 Lazorenko, P.F. 1997, *Kinematics and Phys. Celest. Bodies* 13, 2, 63
 Lazorenko, P.F., & Lazorenko, G.A. 2004, *A&A* 427, 1127
 Lazorenko, P.F. 2006, *A&A* 449, 1271
 Lindegren, L. 1980, *A&A* 89, 41
 Miyamoto, M., & Yoshii, Y. 1995, *AJ* 110, 1427
 Monet, D.G., Dahn, C.C., Vrba, F.J., et al. 1992, *AJ*, 103, 638
 Motch, C., Zavlin, V.E., & Haberl, F. 2003, *A&A*, 408, 323
 Paczyński, B. 1998, *ApJ*, 494, L23
 Pravdo, S., & Shaklan, S. 1996, *AJ* 465, 264
 Pravdo, S.H, Shaklan, S.B., Lloyd, J. et al. 2005, *ASP Conf. Series*, 338, 288
 Pravdo, S.H, Shaklan, S.B., Wiktorowicz, S.J. et al. 2006, *ApJ*, 649, 389
 Quirrenbach, A., Henning, T., & Queloz, D., 2004, *Proc. SPIE* 5491, 424
 Safizadeh, N., Dalal, N., & Griest, K. 1999, *ApJ*, 522, 512
 Udalski, A., Jaroszynski, M., Paczyński, B., et al. 2005, *ApJ*, 628, L109
 Wittkowski, M., Ballester, P., Canavan, T. et al. 2004, *Proc. SPIE* 5491, 617